

# SUSTANE 12-12-12

## 90-Day Controlled Release Fertilizer

Designed for Professional Nursery, Greenhouse & Landscape

### Guaranteed Analysis

Total Nitrogen (N)	12%
5.3% Urea Nitrogen*	
2.9% Nitrate Nitrogen*	
2.3% Ammoniacal Nitrogen*	
0.2% Slowly Available Water Soluble Nitrogen	
1.3% Water Insoluble Organic Nitrogen**	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	12%
Soluble Potash (K <sub>2</sub> O)	12%
Calcium (Ca)	2.00%
Magnesium (Mg)	0.50%
Sulfur (S combined)	1.50%
Iron (Fe)	0.22%
Manganese (Mn)	0.05%
Copper (Cu)	0.01%
Molybdenum (Mo)	0.03%
Boron (B)	0.02%

Derived from aerobically composted turkey litter, hydrolyzed feather meal, sulfate of potash, polymer coated mono-ammonium phosphate, polymer coated potassium nitrate and polymer coated urea.

\*10.3% controlled release nitrogen derived from polymer coated urea, polymer coated potassium nitrate and polymer coated mono-ammonium phosphate.

\*\*1.3% slow release nitrogen derived from aerobically composted turkey litter and hydrolyzed feather meal.

Store in a cool dry place. Keep out of reach of pets and children.

042509

F689

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

### Product Description:

Sustane 12-12-12 was developed as a combination fertilizer to provide all floral and nursery stock with a complete, balanced and controlled release nutrient package. Sustane 12-12-12 provides plant essential nutrients in both controlled release and slow release organic form by combining coated CRF with nutrient rich aerobically composted turkey litter. The Sustane organic component undergoes a 26-week aerobic composting process ensuring that the finished product is stabilized yet biologically active. The controlled release portion of the material is made from industry leading polymer coated urea, polymer coated mono-ammonium phosphate and polymer coated potassium nitrate. Nutrient release rates from Sustane 12-12-12 are based on soil or substrate temperature and microbial activity. The sophisticated two mechanism release matches plant nutrient requirements over the production cycle and prevents dumping of nutrients.

### Direction for Use:

Sustane 12-12-12 provides constant release of nutrients for approximately 90 days when used with container substrates with average temperature of 70°F / 21°C. When determining application rates consider all cultural practices including irrigation, container substrate physical characteristics and growing environment.

**Low Rate:** Recommended for use on salt sensitive species; when concurrent with liquid feed; with production systems utilizing low leaching fractions (less than 10%); and when applied to heavy potting substrates (total porosity less than 65%).

**Medium Rate:** Recommended for use on medium feeding species; on most nursery stock and foliage plants; and when liquid feed is not performed.

**High Rate:** Recommended for use on heavy feeding species; with production systems utilizing high leaching fractions (greater than 20%); and when applied to light, porous substrates (total porosity greater than 80%).

### Container Topdressing Application Rates in grams

Rate	Container Size								
	5" Std.	6" Std.	1 gallon	2 gallon	3 gallon	5 gallon	7 gallon	10 gallon	15 gallon
Low	3	4	6	13	23	37	53	64	69
Med	6	10	15	33	56	90	117	155	167
High	8	15	23	49	86	138	200	238	255

- These rates are intended as guidelines. Sustane encourages a trial prior to changing any fertilizer program.
- Do not store container mix more than two weeks after incorporating fertilizer. Plant material can be damaged from salt accumulation.

### Container Mix & Landscape Application Rates

Container Incorporation Rates by Volume					
lb. per cubic yard			kg. per cubic meter		
Low	Med	High	Low	Med	High
4.5	8.5	14	2.6	5	8.3

Landscape Application Rates by Area					
lb. per 1,000 square feet			kg. per 100 square meter		
Low	Med	High	Low	Med	High
8	16	25	3.9	7.8	12.2

### Bulk Density of Sustane 12-12-12

Rounded Measure (volume)	Grams (weight)
1 teaspoon (tsp)	3.5
1 Tablespoon (Tbs)	10.5
1 oz.	21
1/4 cup	42
1/2 cup	84
1 cup	168

- If mix is stored longer than two weeks, leaching container mix may be required to remove accumulated salts
- Do not steam sterilize container mix after fertilizer has been incorporated
- Monitor electrical conductivity (EC) of container substrate throughout production cycle. Adjust application rate as needed.

# Net Wt. 50 lb. (22.67 kg.)



Sustane Natural Fertilizer, Inc.  
310 Holiday Avenue • PO Box 19  
Cannon Falls, MN 55009 USA  
+1 507-263-3003 • [www.sustane.com](http://www.sustane.com)

