

MEDIUM GRADE



4•6•4 | ALL NATURAL SLOW RELEASE NITROGEN FERTILIZER

SUSTANE
Naturally...



MEDIUM GRADE

Net Weight 50 lb (22.67 kg)

www.sustane.com

4•6•4 | ALL NATURAL SLOW RELEASE NITROGEN FERTILIZER



MEDIUM GRADE



4 • 6 • 4

SUSTÂNE:

Since 1987, Sustâne has been proud to be a part of the global movement toward a more sustainable world: better land management; protection of our waters; saving and rebuilding soils; improving crop yields; helping professional growers, home gardeners, and communities alike - all through the utilization of recycled, renewable, natural resources.

Sustâne Natural Fertilizers are supported by over 25 years of independent research. This is why professional growers worldwide trust in Sustâne for the production of certified organic crops, establishment and maintenance of high quality sports fields, help in stabilizing soils with new vegetation in environmentally sensitive areas, to produce beautiful vegetable gardens and ornamental landscapes and establish and maintain high quality turfgrass.

Sustâne 4-6-4 is a slow-release nitrogen fertilizer made from biologically stable compost, natural sulfate of potash and feather meal.

This product is recommended for use on vegetables, fruits, flowers and landscape beds, tree and shrub establishment, in potting soil mixes and for use as a turfgrass starter fertilizer.

Turf Establishment:

Pre-plant: Broadcast and incorporate into top 4 in (5 cm) of soil.

50 lb per 1,000 ft² or 2,200 lb per acre.
250 g per m² or 2,500 kg per hectare.

Post-plant (45-60 days): Broadcast and water in 25 lb per 1,000 ft² or 1,100 lb per acre.
125 g per m² or 1,250 kg per hectare.

Overseeding:

Apply 25 lb per 1,000 ft² or 125 g per m²

Hydroseeding:

Apply 500 to 4,000 lb per acre (560 to 4,480 kg per hectare).

NOTE: Adjust fertilizer program for local conditions and requirements.

USGA Greens Construction

Sustâne 4-6-4 - 2 mm particle size meets USGA Specifications in combination with sand and peat*

Volume	lb / Yd³ of mix	lb / Ton of mix
1.5%	15.4	11.4

Volume	kg / m³ of mix	kg / Tonne of mix
1.5%	9.1	5.7

ASSUMPTIONS: Bulk density of Sustâne 4-6-4 is 38 lb/cu ft (0.61 g per cm³)

*Providing the sand and peat sources have been approved

Greens Topdressing:

Application Rates (lb per ton) and corresponding amount of Nitrogen applied (lb of N per 1,000 ft²)					
	4-6-4	Core Aeration	Frequent Top Dress		
Volume	Rate lb/ton	0.34 ton/M	1.35 ton/M	0.07 ton/M	0.27 ton/M
1.5%	11.4	0.16	0.62	0.03	0.12

*M = lb per 1,000 ft²

Application Rates (kg per tonne) and corresponding amount of Nitrogen applied (g of N per M²)					
	4-6-4	Core Aeration	Frequent Top Dress		
Volume	Rate kg/ton	3.3 kg/m²	13.2 kg/m²	0.68 kg/m²	2.6 kg/m²
1.5%	5.7	0.62	2.48	0.26	0.49

General Coverage:

50 lb covers 2,000 ft² @ 1 lb N per 1,000 ft².
1,100 lb per acre provides 44 lb N per acre.
22.67 kg covers 186 m² and 125 g covers 1 m²@ 0.5 kg N per 100 m².
1,250 kg per hectare provides 50 kg N per hectare.

Spreader Settings (walk behind)*:

Scotts LawnPro – 15**

Scotts Professional – S**

Scotts Accupro 2000 – M ½***

*Make two passes at each setting. Each pass is a 5 ft (1.5 m) wide interval.

**For 25 lb per 1,000 ft² (125 g per m²)

***For 8.2 lb per 1,000 ft² (40 g per m²)

*Spreader Settings: may vary due to walking speed and ground conditions

Establecimiento de cesped:

Antes de plantar: Esparcir e incorporar en las 4 pulgadas (5 cm), de la parte superior del suelo.

50 libras por 1,000 pies² o 2,200 libras por acre.

250 g por m² o 2,500 kg por hectárea.

Después de plantar: Colocar otra capa, una vez, cada 30-45 días. Esparcir y regar en 25 libras por 1,000 pies² o 1,100 libras por acre (125 g por m² o 1,250 kg por hectárea).

Esparcir la semilla una vez más:

Colocar 25 libras por 1,000 pies²

o 125 g por m².

Esparcir la semilla con agua: Colocar de 500 a 4,000 libras por acre o 560 a 4,480 kg por hectárea.

NOTA: Adaptar el programa de fertilizante para las condiciones y requerimientos locales.

General Guidelines for Garden, Nursery and Landscape Applications

Low Rate: Recommended for salt sensitive species or when applied to heavy, silty clay loam soils or heavy potting substrates (total porosity less than 65%).

Medium Rate: Recommended for use on medium feeding species or when applied to medium, clay loam soils and most potting substrates (total porosity 65 to 80%).

High Rate: Recommended for use on heavy feeding species or when applied to light, sandy soil or light, porous substrates (total porosity greater than 80%).

Container Topdressing Application Rates in grams

Rate	Container Size				
	5" Std.	6" Std.	1 gallon	2 gallon	3 gallon
Low	6	12	14	30	56
Med	14	26	38	78	134
High	20	36	56	118	208

Greenhouse & Nursery Container Mix Rates

Container Mix Rates by Volume					
lb. per cubic yard			kg per cubic meter		
Low	Med	High	Low	Med	High
11	20	30	6.5	11.8	17.8

Landscape & Flower Application Rates by Area					
lb. per 1,000 square feet			kg per 100 square meter		
Low	Med	High	Low	Med	High
18	36	56	8.8	17.6	27.4

Seed Starting/Bedding Plants Rates by Volume

Mix Rates by Volume					
lb. per cubic yard			kg per cubic meter		
Low	Med	High	Low	Med	High
6	9	12	3.6	5.3	7.1

Vegetable Gardens Application Rates by Area*

Plant	Tbsp / ft²	Lb / 100 ft²	Grams / M²
Tomatoes	2.75	5.75	281
Lettuce	2.5	5	244
Cucumbers	1.33	2.75	134
Peppers	0.4	0.8	39

*Visit www.sustane.com for rates for other specific crops.

*Adjust to higher rates for areas with higher plant densities.

Amending Landscape Soil Beds:

Scratch or work Sustâne into the top 2 to 4 in (5 to 10 cm) of the soil prior to planting. Reapply at 45 days, spreading evenly over the bed surface.

Reparación de una extensión del terreno: Preparar o colocar Sustane en la superficie de 2 a 4 pulgadas (5 a 10 cm) del suelo antes de plantar. Repetir a los 45 días, esparciendo uniformemente sobre la superficie de la extensión del suelo a trabajar.

Fertilizing Shrubs and Trees:

Apply ¼ cup of Sustane for plants with a 2 ft. diameter canopy (60 cm² for 60 cm diameter canopy). Increase rates by ¼ cup for every additional 6 in. of canopy spread (60 cm² per 15 cm). Apply at planting or in the spring and fall for established trees and shrubs.

Fertilización de árboles y arbustos: Colocar ¼ de taza de Sustane para plantas que tienen una copa de 2 pies de diametro y de ancho (60 cm³ por 60 cm de diametro). Aumentar las proporciones, con ¼ de taza, por cada 6 pulgadas más del tamaño de la copa (60 cm³ por 15 cm). Colocar fertilizante al plantar en la primavera, y en el otoño para los árboles y arbustos establecidos.

Tree Planting Rate Per Tree

	Tree trunk caliper (in)¹				
	1.25" (3cm)	2" (5cm)	3" (7.5cm)	4" (10cm)	5" (12.7cm)
rate (lb)	0.67	1.0	2.0	3.5	5
rate (kg)	0.3	0.45	0.9	1.5	2.0

¹ Tree trunk caliper is measured 6 in (15 cm) above the soil surface

Tree Maintenance Rate Per Tree

	Tree trunk caliper (in)¹				
	1.25" (3cm)	2" (5cm)	3" (7.5cm)	4" (10cm)	5" (12.7cm)
rate (lb)	1.33	1.5	2.33	4.33	6
rate (kg)	0.6	0.45	0.9	1.5	2.5

Agricultural Crops:

**Recommended fertilizer application rates for agricultural crops are crop, crop yield goal, soil and climate specific. Please refer to your soil test results or to the recommendations of your area agricultural extension agent and/or crop consultant for specific recommendations. Additional crop specific application information and fertilizer research is available at www.sustane.com. You can also contact your regional Sustane technical representative for additional guidance.

Productos Agrícolas:

****Las proporciones para colocar el fertilizante recomendadas para los productos agrícolas cosechados son el rendimiento máximo de cocehas específico para el clima y suelo del lugar.**

Guaranteed Analysis:

Total Nitrogen (N). 4%

0.4% Ammoniacal Nitrogen

0.4% Other Water Soluble Nitrogen

3.2% Water Insoluble Nitrogen*

Available Phosphate (P₂O₅). 6%

Soluble Potash (K₂O). 4%

Calcium (Ca). 4%

Magnesium (Mg). 1.00%

Sulfur (S). 1.00%

Iron (Fe). 0.50%

Manganese (Mn) 0.05%

Derived from aerobically composted turkey litter, feather meal and sulfate of potash.

*3.2% slowly available nitrogen from aerobically composted turkey litter and feather meal

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

Store in a cool, dry place.

Keep out of reach of pets and children

Almacenelo en un lugar oscuro, fresco y seco.

Mantengalo fuera del alcance de los niños y mascotas.

NET WEIGHT 50 IB (22.7 kg)

PESO NETO 50 IB (22.7 kg)

Granulation Specifications:

Medium Grade: Mesh Size -7+14
(2.8 mm to 1.4 mm) SGN 200

F689

Manufactured by Sustâne® Natural Fertilizer, Inc.
310 Holiday Avenue • Cannon Falls, MN 55009 U.S.A.
+1(507) 263-3003 • e-mail: help@sustane.com

www.sustane.com



Visit www.sustane.com for more information.



Notice: Read Before Using Product
Use of this product is subject to the following disclaimer of warranties and conditions (including limitations of liability).
Disclaimer of Warranties: There are no warranties, express or implied, or merchantability or of fitness for a particular purpose or otherwise, which extend beyond the printed statements on this package. Under no circumstances shall Sustâne® Natural Fertilizer, Inc. be liable for any incidental or consequential damage resulting from a failure to use this product in accordance with the printed directions on this package.
Conditions: The printed directions on this package are for the protection of the user. Any use of this product not in conformity with those printed directions is at the risk of the user.
Limitations of Liability: In no event shall Sustâne® Natural Fertilizer, Inc.'s liability for damages arising out of the use of this product exceed the purchase price of the product.

